

February 2, 2008

COSMIC MICROWAVE BACKGROUND TEMPERATURE AT GALAXY CLUSTERS

E.S. Battistelli, M. DePetrìs, L. Lamagna, F. Melchiorri, E. Palladino, G. Savini

Department of Physics, University "La Sapienza", P.le A. Moro 2, 00185, Rome, Italy

A. Cooray

Theoretical Astrophysics, California Institute of Technology, Pasadena CA 91104

A. Melchiorri

University of Oxford, Denys Wilkinson building, Astrophysics, Keble Road, Oxford, OX1-3RH, UK

Y. Rephaeli, M. Shimon

School of Physics and Astronomy, Tel Aviv University

ABSTRACT

We have deduced the cosmic microwave background (CMB) temperature in the Coma cluster (A1656, $z = 0.0231$), and in A2163 ($z = 0.203$) from spectral measurements of the Sunyaev-Zel'dovich (SZ) effect over four passbands at radio and microwave frequencies. The resulting temperatures at these redshifts are $T_{Coma} = 2.789^{+0.080}_{-0.065}$ K and $T_{A2163} = 3.377^{+0.101}_{-0.102}$ K, respectively. These values confirm the expected relation $T(z) = T_0(1+z)$, where $T_0 = 2.725 \pm 0.002$ K is the value measured by the COBE/FIRAS experiment. Alternative scaling relations that are conjectured in non-standard cosmologies can be constrained by the data; for example, if $T(z) = T_0(1+z)^{1-a}$ or $T(z) = T_0[1 + (1+d)z]$, then $a = -0.16^{+0.34}_{-0.32}$ and $d = 0.17 \pm 0.36$ (at 95% confidence). We briefly discuss future prospects for more precise SZ measurements of $T(z)$ at higher redshifts.

Subject headings: cosmic microwave background—cosmology:observations—galaxies:clusters:individual(A1656,A2163)

1. Introduction

The (present) CMB temperature was precisely measured by the Far Infrared Absolute Spectrometer (FIRAS) on board the COBE satellite, $T_0 = 2.725 \pm 0.002$ K, in the frequency range $2\text{--}20\text{ cm}^{-1}$ (Mather *et al.* 2001). These measurements essentially rule out all cosmological models in which the CMB spectrum is non-Planckian at $z = 0$. Models with a purely blackbody spectrum but with a different $T(z)$ dependence than in the standard model are, however, unconstrained by the FIRAS database. Also unconstrained are models with spectral distortions that are now negligible, but were appreciable in the past. A specific example is the relation $T(z) = T_0(1+z)^{1-a}$, where a is a parameter of the theory (see, e.g., Lima *et al.* 2000). More generally, models in which ratios of some of the fundamental constants vary over cosmological time are also of considerable interest.

So far, $T(z)$ has been determined mainly from measurements of microwave transitions in interstellar clouds which contain atoms and molecules that are excited by the CMB (as reviewed by LoSecco *et al.* 2001). The temperature has been determined in the Galaxy, as well as in clouds at redshifts up to $z \sim 3$ (Levshakov *et al.* 2001). These measurements are affected by substantial systematic uncertainties stemming from the unknown physical conditions in the absorbing clouds (Combes and Wiklind 1999).

The possibility of determining $T(z)$ from measurements of the Sunyaev-Zel'dovich (SZ) effect had been suggested long ago (Fabbri, Melchiorri & Natale 1978, Rephaeli 1980). (For general reviews of the effect and its cosmological significance, see Rephaeli 1995a, Birkinshaw 1999.) The proposed method is based on the steep frequency dependence of the change in the CMB spectral intensity, ΔI , due to the effect, and the weak dependence of ratios $\Delta I(\nu_i)/\Delta I(\nu_j)$ of intensity changes measured at two frequencies (ν_i, ν_j) on properties of the cluster (Rephaeli 1980). Because of this, and the fact that – in the standard cosmological model – the effect is essentially independent of z , SZ measurements have the potential of yielding much more precise values of $T(z)$ than can be obtained from ratios of atomic and molecular lines. With the improved capability of reasonably precise spectral measurements of the SZ effect, the method can now be used to measure $T(z)$ in nearby and moderately distant clusters. Here we report first results from spectral analysis of SZ measurements in the Coma and A2163 clusters of galaxies.

2. $T(z)$ from SZ

The CMB intensity change due to Compton scattering in a cluster can be written in the form

$$\Delta I = \frac{2k^3 T^3}{h^2 c^2} \frac{x^4 e^x}{(e^x - 1)^2} \int d\tau \left[\theta f_1(x) - \beta + R(x, \theta, \beta) \right], \quad (1)$$

where $x = h\nu/kT$ is the non-dimensional frequency, $\theta = kT_e/mc^2$ is the electron temperature in units of the electron rest energy, and β is the line of sight (los) component of the cluster (peculiar) velocity in the CMB frame in units of c . The integral is over the Compton optical depth, τ . Both the thermal (Sunyaev & Zeldovich 1972) and kinematic (Sunyaev & Zel'dovich 1980) components of the effect are included in equation (1), separately in the first two (additive) terms, and jointly in the function $R(x, \theta, \beta)$. In the non-relativistic limit (which is valid only at low electron temperatures and frequencies) the spectral dependence of ΔI is fully contained in the product of the x -dependent pre-factor times the function $f_1(x) = x(e^x + 1)/(e^x - 1) - 4$. The more exact treatment of Compton scattering in clusters necessitates a relativistic calculation (Rephaeli 1995b) due to the high electron velocities. The function $R(x, \theta, \beta)$ includes the additional spectral, temperature, and (cluster) velocity dependence that is obtained in a relativistic treatment. This function can be approximated by an analytic expression that includes terms to orders θ^5 and $\beta^2\theta$:

$$R(x, \theta, \beta) \simeq \theta^2 \left[f_2(x) + \theta f_3(x) + \theta^2 f_4(x) + \theta^3 f_5(x) \right] - \beta \theta \left[g_1(x) + \theta g_2(x) \right] + \beta^2 \left[1 + \theta g_3(x) \right]. \quad (2)$$

The spectral functions f_i and g_i were determined by Itoh *et al.* (1998), Itoh *et al.* (2002), and Shimon & Rephaeli (2002). For our purposes here this analytic approximation is sufficiently exact even close to the crossover frequency. The non-relativistic limit, $R \equiv 0$, applies if the sum of all these terms can be ignored at the desired level of accuracy.

The z dependence of ΔI is fully determined by the functions $\nu = \nu(z)$, and $T = T(z)$. The temperature-redshift relation may assume various forms in non-standard cosmologies; here we consider two examples. In the first, $T(z) = T(0)(1+z)^{(1-a)}$, where a is taken to be a free parameter, but with the standard scaling $\nu = \nu_0(1+z)$ unchanged. With these relations the non-dimensional frequency obviously depends on z , $x = x_0(1+z)^a$, if $a \neq 0$; here, $x_0 = h\nu_0/kT(0)$. Another functional form which seems also to be of some theoretical interest is $T(z) = T(0)[1+(1+d)z]$ (LoSecco *et al.* 2001), for which $x = x_0(1+z)/[1+(1+d)z]$. Obviously, in the standard model $a = d = 0$.

For a slow moving ($\beta < 10^{-3}$) cluster, the expression for ΔI in the non-relativistic limit depends linearly on the Comptonization parameter, $y = \int \theta d\tau$, which includes all dependence on the cluster properties. A ratio of values of ΔI at two frequencies is then essentially independent of these cluster properties. In the more general case, the first term

in the square parentheses in eq. (1) still dominates over the other two, except near the crossover frequency (whose value generally depends on T_e , except in the non-relativistic limit where $x_c = 3.83$; Rephaeli 1995b, Nozawa *et al.* 1998, Shimon & Rephaeli 2002), where the sum of the temperature dependent terms vanish. For values of x outside some range (roughly, $3.5 < x < 4.5$), the dependence of ΔI , and – particularly, a ratio of values of ΔI – on β is very weak since the observed temperature range in clusters corresponds to $0.006 < \theta < 0.03$, whereas typically $\beta < 0.002$.

3. Data Analysis

We have analyzed results of SZ measurements in the Coma cluster (A1656) and A2163. Measurements of Coma, $z = 0.0231 \pm 0.0017$ (Struble and Rood 1999) were made with the MITO (DePetrìs *et al.* 2002) telescope in 20 hours of integration. We also use the result of measurements at 32 GHz made with the OVRO 5.5m telescope (Herbig *et al.* 1995, Mason *et al.* 2001). A2163, $z = 0.203 \pm 0.002$ (Arnaud *et al.* 1992), was observed with the SuZIE array (Holzapfel *et al.* 1997a), and with both the OVRO and BIMA interferometric arrays (LaRoque *et al.* 2000).

When observing a cluster the SZ part of the measured signal is

$$\Delta S_i = G_i A\Omega|_i \int_0^\infty \Delta I(0) \epsilon_i(\nu) d\nu, \quad (3)$$

where G_i is the responsivity of the i^{th} photometric channel, $A\Omega|_i$ is the corresponding throughput, and $\epsilon_i(\nu)$ is the spectral efficiency. The full measured signal includes also contributions from the atmosphere, CMB anisotropies, and – at very high frequencies – also emission from dust. Multifrequency observations allow us to remove contributions from both the primary CMB anisotropy and the kinematic SZ effect, as has been attempted in the analysis of MITO measurements of the Coma cluster (DePetrìs *et al.* 2002).

The ratio of signals in two different photometric channels i and j is

$$\frac{\Delta S_i}{\Delta S_j} = \frac{G_i A\Omega|_i}{G_j A\Omega|_j} \frac{\int_0^\infty \frac{x^4 e^x}{(e^x - 1)^2} \left\{ \int d\tau \left[\theta f_1(x) - \beta + R(x, \theta, \beta) \right] \right\} \epsilon_i(\nu) d\nu}{\int_0^\infty \frac{x^4 e^x}{(e^x - 1)^2} \left\{ \int d\tau \left[\theta f_1(x) - \beta + R(x, \theta, \beta) \right] \right\} \epsilon_j(\nu) d\nu}. \quad (4)$$

The main dependence on the cluster properties in y cancels out when β is negligible. Multifrequency observations that include measurements at the crossover frequency (e.g., MITO) afford effective separation of the thermal and kinematic components, exploiting their very different spectral shapes. This was demonstrated in the analysis of MITO measurements

of the Coma cluster (De Petris et al. 2002). Since the above ratio depends weakly on the cluster velocity, the residual uncertainty due to velocities of even ± 500 km/s can be ignored in comparison with other errors. The ratio is also weakly dependent on the gas temperature at a level which we found to correspond to $\sim 1\%$ uncertainty in the estimation of the CMB temperature (for a typical observational error in T_e). Moreover, the uncertainty associated with the absolute calibration, G_i , is largely removed once we fit data from several photometric channels, as long as they are calibrated with a source with a known spectrum (e.g., a planet) even if its absolute calibration is uncertain. Only relative uncertainties among the various spectral channels are important; these include differences in angular response and in atmospheric transmittance. A standard blackbody source with a precisely calibrated temperature is therefore not required. We expect these considerations to imply that the precision of CMB temperature measurements via the SZ effect will not be appreciably affected by most of the known systematic errors. The level of precision in the measurement of $T(z)$ is limited largely by other observational uncertainties, as discussed below.

The responsivity G_i of each channel is usually determined from detailed observations of very well measured sources such as planets (mostly Jupiter or Saturn) and the Moon. While the temperature uncertainty of these sources can be as large as 10%, their spectra are relatively well known. For a source at a temperature of T_S and with a throughput Ω_S , the signal in the Rayleigh-Jeans part of the spectrum is

$$\Delta S_i = G_i A_i \Omega_S T_S \frac{2k}{c^2} \int_0^\infty \nu^2 \epsilon_i(\nu) \eta(\nu) d\nu. \quad (5)$$

Since we are interested in the ratio of the signals in two channels, T_S drops out from the final expression, and the uncertainty in the ratio of values of the brightness temperature in the two channels depends only on the emissivity of the planet, $\eta(\nu)$, convolved with the spectral efficiencies of the channels, $\epsilon_i(\nu)$. Note that the throughputs of individual channels, $A\Omega|_i$, are usually made almost equal by an appropriate choice of the optical layout.

The main uncertainty in the MITO measurements is due to imprecise knowledge of the bandwidth and transmittance of the filters. Therefore, we have precisely measured the total efficiency of our photometer by means of a lamellar grating interferometer. The response of the photometer is measured when it is illuminated with a laboratory blackbody at different temperatures ranging from liquid N_2 to room temperature. The effect of the uncertainty in our bandwidths has two main consequences. The first is the error in the ratio of responsivities of channel i and channel j , G_i/G_j . We estimate this error by computing the ratio of expected signals evaluated by convolving Moon and Jupiter spectra with the spectra of our filters, and taking into account spectroscopic uncertainties; we obtain a final error estimate which is less than 1%. The second consequence of bandwidth uncertainty enters in the integrations over

the SZ spectral functions in eq.[4]. However, this latter error is smaller than the former and is practically negligible.

Also quite precisely measured were the angular responses of the telescope and the photometer in the four channels. We have studied differences in the response of the four channels to both a point source, such as Jupiter, and an extended source, such as the Moon, as they cross the field of view. The optical layout of our photometer is such that all four detectors observe the same sky region. The differences among channels are less than 1% for extended sources. Finally, we have also studied the change in relative efficiency of our observations under different atmospheric conditions. The simulations show a change of ~ 1 mK in the estimated CMB temperature if the water vapor content changes from 0.5 mm to 1.5 mm, as determined by convolving theoretical atmospheric spectra with our filters. The uncertainties in determining the water vapor content, related to our procedure of measuring the atmospheric transmittance through a sky-dip technique, are at a level of 30%. However, when the effect of this error is propagated through our procedure and analysis we estimate that it amounts to a negligible error in the final result for the CMB temperature.

In conclusion, systematic errors contribute less than 3% to the observed MITO signals, and are quite negligible with respect to the 10% contribution associated with the residual noise of the four detectors. For the OVRO measurement, we use the total error as specified by Mason *et al.* (2001); the statistical weight of this low frequency measurement is such that it only contributes $\simeq 12\%$ to the final results.

Our assessment of the errors in the SuZIE and OVRO & BIMA measurements of A2163 (LaRoque *et al.* 2000) is far less certain than those in the MITO measurements, since we do not know the exact spectral responses of their filters and the atmospheric conditions during each of these observations. We have used Gaussian profiles for the SuZIE filters, with peak frequencies and FWHM values as given by Holzapfel *et al.* (1997a). For the OVRO & BIMA data point, we also take a Gaussian profile with a peak frequency at 30 GHz and a 1 GHz bandwidth. We have employed data provided by SuZIE, for which a correction due to thermal dust emission has been taken into account. The Coma and A2163 data are collected in Table 1 and Table 2. Finally, the value we used for the electron temperature is (8.25 ± 0.10) keV, as measured with XMM in the central region of Coma (Arnaud *et al.* 2001); the value for A2163 is $12.4^{+2.8}_{-1.9}$ keV (Holzapfel *et al.* 1997b).

In both sets of cluster observations, we have minimized the difference between theoretical and experimental ratios (properly weighted by statistical errors) with T as a free parameter. The results are $T_{Coma}(0) = 2.726^{+0.078}_{-0.064}$ K which, when multiplied by $(1+z)$ with $z = 0.0231 \pm 0.0017$, yields

$$T_{Coma} = 2.789^{+0.080}_{-0.065} \text{K}. \quad (6)$$

The main uncertainty is in the SZ observations, with only a small error (7×10^{-4} K) due to the uncertainty (0.1 keV) in the electron temperature. For A2163 we obtain $T_{A2163}(0) = 2.807^{+0.084}_{-0.085}$ K, and when multiplied by $(1+z)$ with $z = 0.203 \pm 0.002$, yields

$$T_{A2163} = 3.377^{+0.101}_{-0.102} \text{ K}. \quad (7)$$

The larger error in T_e (~ 2 keV) translates to an uncertainty of 1.5×10^{-2} K in the deduced CMB temperature at the redshift of A2163. The mean of the above two values for $T(0)$ is $\langle T \rangle = 2.766^{+0.058}_{-0.053}$ K, in good agreement with the COBE value.

Using the COBE value of the temperature and the two values deduced here for $T(z)$ at the redshifts of Coma and A2163, we fit the three data points by the relation $T(z) = T(0)(1+z)^{1-a}$; the best fit is shown by the dashed line in Figure 1. It corresponds to $a = -0.16^{+0.34}_{-0.32}$ at 95% confidence level (CL). With the alternative scaling $T(z) = T(0)[1 + (1+d)z]$, we obtain a best fit value of $d = 0.17 \pm 0.36$ at the 95% CL.

These first SZ results for $T(z)$ are clearly consistent with the standard relation, although the most probable values of both a and d indicate a slightly stronger z dependence. Our values for these two parameters are quite similar to those deduced by LoSecco & Mathews (2001) from measurements of microwave transitions. Their results – based on two firm detections at redshifts 2.338 and 3.025 – are $a = -0.05 \pm 0.13$ and $d = 0.10 \pm 0.28$ (at 95% CL). Thus, our SZ-based values already provide the same level of precision even though the two clusters are at much lower redshifts ($z \lesssim 0.2$). SZ measurements of higher redshift clusters will clearly provide a preferred alternative to the atomic and molecular lines method. If we fit the above relations to the combined data sets (consisting of all five temperatures), we obtain $a = -0.09 \pm 0.20$ and $d = 0.14 \pm 0.28$ (at 95% CL). Finally, Molaro *et al.* (2002) have recently questioned the validity of the temperature measurement at $z = 2.338$ (by LoSecco & Mathews 2001), and have revised the value at $z = 3.025$ to $12.1^{+1.7}_{-3.2}$ K. We have thus repeated the fit excluding the fourth data point and using the latter value for the fifth; the results from this fit are essentially unchanged: $a = -0.11 \pm 0.22$, and $d = 0.16 \pm 0.32$.

4. Discussion

The method employed here to measure $T(z)$ can potentially yield very precise values which will tightly constrain alternative models for the functional scaling of the CMB temperature with redshift, and thereby provide a strong test of non-standard cosmological models. In order to improve the results reported here, multi-frequency measurements of the SZ effect in a significantly larger number (~ 20) of clusters are needed. The clusters should include nearby ones in order to better understand and control systematic errors. Exactly such ob-

servations are planned for the next 2-3 years with the upgraded MITO project (Lamagna *et al.* 2002), and the stratospheric BOOST experiment (Lubin P., private communication) and OLIMPO experiment (Masi *et al.* 2003). These experiments will employ sensitive bolometric arrays at four frequency bands with high spatial resolution.

As detector noise is reduced, the uncertainty in the electron temperature may contribute the dominant relative error. For example, an uncertainty of ~ 2 keV – as in the case of Abell 2163 – becomes dominant when detector noise is reduced to a level ten times lower than the values used here. Therefore, it will be necessary to select clusters for which the gas temperature is precisely (to a level of ~ 0.1 keV) known. With the large number of clusters that have already been – or will be – sensitively observed by the XMM and *Chandra* satellites, it should be possible to optimally select the SZ cluster sample.

With the currently achievable level of precision in intracluster gas temperature measurements, the present technique makes it possible to determine a and d with an uncertainty that is not less than 0.03 on both parameters, even with ideal (i.e., extremely high sensitivity) SZ measurements. The projected sensitivities of the future Planck and Herschel satellites will enable reduction of the overall error in the values of these two parameters by more than an order of magnitude. In addition to limits on alternative CMB temperature evolution models, such a level of precision will open new possibilities of testing the time variability of physical constants. For example, it will be feasible to measure a possible variation of the fine structure constant, thereby providing an alternative to the current principal method which is based on spectroscopic measurements of fine structure lines in quasar absorption spectra.

REFERENCES

- Arnaud M., Aghanim N., Gastaud R., Neumann D. M., Lumb D., Briel U., Altieri B., Ghizzardi S., Mittaz J., Sasseen T. P. and Vestrand W. T., 2001, A&A 365, L67-L73
- Arnaud M., Hughes J. P., Forman W., Jones C., Lachize-Rey M., Yamashita K. and Hatsuakade I., 1992, ApJ 390:345-349
- Birkinshaw M., 1999, Phys. Rep. 310, 97
- Combes F. and Wiklind T., 1999, ASP Conf. Ser. 156: Highly Redshifted Radio Lines, 210, ed. by Carilli C. L., Radford S. J. E., Meten K. M. & Langston G. I., San Francisco: ASP (1999), Proc. Conf. Green Bank, WV, USA, 9-11 October
- De Petris M., D’Alba L., Lamagna L., Melchiorri F., Orlando A., Palladino E., Rephaeli Y., Colafrancesco S., Kreysa E., and Signore M., 2002, ApJ, 574, L119-L122

- Fabbri R. and Melchiorri. F. , & Natale V. 1978, *Astrophysics and Space Science*, 59, 223
- Herbig T., Lawrence C.R. and Readhead A.C.S., 1995, *ApJ* 449, 5
- Holzappel W. L., Wilbanks T. M., Ade P. A. R., Church S. E., Fischer M. L., Mauskopf P. D., Osgood D. E. and Lange A. E., 1997a, *ApJ* 479, 17
- Holzappel W. L., Arnaud M., Ade P. A. R., Church S. E., Fisher M. L., Mauskopf P. D., Rephaeli Y., Wilbanks T. M., & Lange A. E., 1997b, *ApJ* 480, 449
- Itoh N., Kohyama Y. and Nozawa S., 1998, *ApJ* 502, 7
- Itoh N., Sakamoto T., Kusano S., Kawana Y. and Nozawa S., 2002, *A&A* 382, 722-729
- Lamagna L., De Petris M., Melchiorri F., Battistelli E. S., De Grazia M., Luzzi G., Orlando A., Savini G., 2002, *AIP Conference Proceedings*, 616, 92, editors: M. De Petris and M. Gervasi
- LaRoque S. J., Carlstrom J. E., Reese E. D., Holder G. P., Holzappel W. L., Joy M. and Grego L., 2002, *astro-ph/0204134*, submitted to *ApJ*
- Levshakov S.A., *et al.* 2001, in *Proceed. Workshop on Chemical Enrichment of Intracluster and Intergalactic Medium* (May 14-18 2001, Vulcano, Italy), *PASP*, in press
- Lima J. A. S., Silva A. I. and Viegas S.M., 2000, *MNRAS* 312, 747-752
- LoSecco J. M., Mathews G. J. and Wang Y., 2001, *astro-ph/0108260*, submitted to *Phys. Rev. D*
- Masi S., *et al.* 2003, to appear in the proceedings of the 4th National Conference on Infrared Astronomy, 4-7 December 2001, Perugia, eds. S. Ciprini, M. Busso, G. Tosti and P. Persi, "Memorie della Societ Astronomica Italiana", 74, 1
- Mason B.S., Myers S.T. and Readhead A.C.S., 2001, *ApJL* 555, L11
- Mather J. C., Fixsen D. J., Shafer R. A., Mosier C. and Wilkinson D. T., 1999, *ApJ* 512, 511-520
- Molaro P., Levshakov S. A., Dessauges-Zavadsky M. and D'Odorico S., 2002, *A&A* 381, L64-L67
- Nozawa S., Itoh N. and Kohyama Y., 1998, *ApJ* 508, 17
- Rephaeli Y., 1980, *ApJ* 241, 858

Rephaeli Y. 1995a, *Ann. Rev. Astron. & Ap.* 33, 541

Rephaeli Y. 1995b, *ApJ* 445, 33

Shimon M. & Rephaeli Y., 2002, *ApJ* 575, 12S

Struble M. F. and Rood H. J., 1999, *ApJ* 125, 35-71

Sunyaev R.A. & Zel'dovich Ya. B., 1972, *Comm. Astrphys. Space Phys.*, 4, 173

Sunyaev R.A. & Zel'dovich Ya. B., 1980, *MNRAS* **190**, 413




Fig. 1.— Black diamonds mark values of the CMB temperature obtained by COBE/FIRAS and SZ measurements. The two values reported by LoSecco *et al.* (2001) are shown as white squares, and the white diamond is the value reported by Molaro *et al.* (2002). The solid line is $(1+z)$ scaling law as predicted in the standard model. The dashed line shows the best fit to the alternative scaling law with a determined by the COBE/FIRAS and SZ measurements, whereas dot-dashed lines are the $\pm 1\sigma$ values of a . The inset shows a magnified view of the region $z \leq 0.25$.

Table 1. SZ signals from COMA used in the present analysis.

Experiment	Frequency [GHz]	Bandwidth [GHz]	ΔI [$MJsr^{-1}$]	ΔT [μK]
OVRO	32.0	13.0	-0.0159 ± 0.0028	-520 ± 93
MITO1	143	30	-0.068 ± 0.014	-179.3 ± 37.8
MITO2	214	30	-0.016 ± 0.039	-33.4 ± 81.2
MITO3	272	32	0.075 ± 0.015	169.8 ± 35.1

Table 2. SZ measurements of A2163 used in the present analysis.

Experiment	Frequency [GHz]	Bandwidth [GHz]	ΔI [$MJsr^{-1}$]	ΔT [μK]
OVRO+BIMA	30	1	-0.048 ± 0.006	-1777 ± 222
SuZIE1	141.59	12.67	-0.380 ± 0.037	-1011.3 ± 98.0
SuZIE2	216.71	14.96	-0.103 ± 0.077	-213.0 ± 159.3
SuZIE3	268.54	25.66	0.295 ± 0.105	662.2 ± 235.7